

Rx Only
For topical use only

Federal Law restricts this device to sale by or on the order of a licensed healthcare practitioner.

[EN] Description

StrataMark is a rapidly drying, non-sticky, transparent gel formulation for the management of Striae Distensae (stretch marks).

When used as directed, StrataMark dries to form a thin, flexible and protective layer, that is gas permeable and waterproof which hydrates and protects stretch marks, allowing the skin to normalize the collagen synthesis cycle.

StrataMark is suitable for all areas of the skin, including exposed and sensitive areas, as well as large areas of the body (i.e. buttocks, abdomen, thighs, breasts and back).

Indication for use

StrataMark is intended to be used under the direction of healthcare practitioners in the management of stretch marks.

StrataMark is indicated for the management of all types of stretch marks, particularly those covering a wide surface area, resulting from pregnancy, adolescent growth spurts, weight gain or loss, or bodybuilding.

StrataMark is indicated to soften and flatten stretch marks, reduce redness and discoloration associated with stretch marks as well as to relieve the itching and discomfort of stretch marks.

It contains no alcohols, parabens or fragrances.

StrataMark is suitable for pregnant women, breastfeeding mothers, children and people with sensitive skin.

StrataMark is intended for single patient use.

Directions for use



On the first use of 1.75 oz (50 g) tubes remove the cap, then the protective seal and close with the cap after use.

Ensure that the skin is clean and dry.

Apply a very thin layer of StrataMark to the relevant areas with the fingertips and distribute evenly to form a very thin layer and allow the gel to dry.

When applied, StrataMark should be dry in 5–6 minutes.

If it takes longer to dry you have probably applied too much. Gently remove the excess with a clean tissue or gauze and allow the drying process to continue.

Once dry, StrataMark can be covered with undergarments, sunscreens or cosmetics.

StrataMark should be applied once per day or as advised by your physician.

For best results StrataMark should be maintained in continuous contact with the skin (24 hours a day/7 days a week).

ADVANCED FORMULATION

Strata

for the management
of stretch marks

mark[®]

How much StrataMark do I need?

StrataMark gel is an advanced formulation that requires substantially less product per application than typical creams or gels.

StrataMark 1.75 oz (50 g) is enough to treat an area of 12×12 inch (30×30 cm) for 30 days.

Warning

- For external use only.
- StrataMark should not be applied to third degree burns or to open wounds.
- StrataMark should not be placed in contact with eyes.
- StrataMark should not be applied over other skin treatments without the advice of your physician.
- StrataMark may stain clothing if not completely dry. If staining occurs, dry cleaning should be able to remove it without damaging the fabric.
- For correct storage please reclose the tube tightly with the cap.
- If irritation occurs, discontinue use and consult your physician.
- Keep out of the reach of children.
- Do not use after the expiration (EXP) date printed on the tube. The expiration (EXP) date does not change once the tube has been opened.
- Do not use if the tube is damaged.

Contraindications:

Do not administer to patients with known hypersensitivity to the ingredients of this product.

Ingredients: Dimethylpolysiloxane, dihydroxysiloxane and alkylmethylsiloxane.

How supplied:

StrataMark 1.75 oz (50 g) tube

Product identification code:

73661-423-50



Consult instructions for use



Use by date




Batch code



86°F

Upper limit of temperature

 Stratpharma AG
Aeschenvorstadt 57
CH-4051 Basel
Switzerland

Distributed by
Marnel Pharmaceuticals LLC
Charleston, SC 29403, U.S.A.

Date of preparation of this
information: 04/2020

 Stratpharma
Switzerland

 Marnel Pharmaceuticals

us.stratamark.net
1-833-788-0312



SK000USENP02-0420